Government of Saskatchewan

Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 12, July 24, 2014

For the Period July 15 to 21, 2014

One year ago
Livestock producers had 69 per cent of the hay crop cut and 43 per cent baled or put into silage. Crops were in mostly good condition.

Follow the 2014 Crop Report on Twitter @SKAgriculture

Haying continues in the province and livestock producers now have 42 per cent of the hay crop baled or put into silage, with an additional 49 per cent cut and ready for baling, according to Saskatchewan Agriculture's weekly Crop Report.

Rain showers and high humidity have delayed haying and decreased hay quality in some areas. Thirteen per cent is rated as excellent in quality, 77 per cent good, eight per cent fair and two per cent poor. Hay yields are slightly below the five-year average (2009-2013). The estimated average hay yields on dry land are reported as 1.3 tons per acre for alfalfa, 1.4 tons per acre for alfalfa/brome hay, 1.1 tons per acre for other tame hay, 0.9 tons per acre for wild hay and 1.7 tons per acre for greenfeed. On irrigated land, the estimated average hay yields are 1.9 tons per acre for alfalfa, 1.8 tons per acre for alfalfa/brome hay, 1.4 tons per acre for other tame hay and 2.3 tons per acre for wild hay and greenfeed.

Rain during the week ranged from trace amounts to 84 mm in the Nipawin area. Across the province, topsoil moisture on cropland is rated as 14 per cent surplus, 75 per cent adequate, 10 per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 74 per cent adequate, 13 per cent short and three per cent very short. Some areas in the south are drier than normal and will soon need moisture to help crops mature and fill.

Warm weather has helped advance many crops and the majority are in fair to excellent condition.

Storms moved through the province last week, bringing strong winds, heavy rain and hail. Other sources of crop damage include insects such as grasshoppers and wheat midge and diseases such as root rot and leaf spots.

Farmers are busy haying and controlling insects and crop disease.

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Southeastern Saskatchewan (Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas; Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas; Crop District 3ASE – Radville and Lake Alma areas)

Warm weather is helping to advance many crops in the region; however, many areas are in need of rain to help crops mature and fill. Rainfall varied across the region, ranging from trace amounts to 37 mm in the Moose Jaw area. The Moosomin area has received 600 mm of rain since April 1, the greatest amount for both the region and the province.

Estimated Southeast Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.3	N/A
Brome/Alfalfa	1.5	N/A
Other Tame Hay	1.2	N/A
Wild Hay	1.0	N/A
Greenfeed	1.7	N/A

Topsoil moisture conditions on cropland are rated as 15 per cent surplus, 72 per cent adequate and 13 per cent short. Hay land and pasture topsoil moisture is rated as 13 per cent surplus, 71 per cent adequate and 16 per cent short. Moisture conditions vary across the region. CD 1B is reporting that 29 per cent of the cropland has surplus topsoil moisture, while CDs 2A and 3ASE are reporting that 50 per cent of the cropland there is short topsoil moisture.

Livestock producers now have 37 per cent of the hay crop baled or put into silage, with an additional 23 per cent cut and ready for baling. Hay quality is rated as 11 per cent excellent, 82 per cent good, four per cent fair and three per cent poor.

The majority of crop damage is due to localized flooding, wind, hail and drought conditions. Producers are spraying for insects such as wheat midge and grasshoppers, and for diseases such as fusarium head blight, sclerotinia and leaf spots.

Farmers are busy haying, controlling pests and hauling grain.

Southwestern Saskatchewan (Crop District 3ASW – Coronach, Assiniboia and Ogema areas; Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas; Crop District 3B – Kyle, Swift Current , Shaunavon and Ponteix areas; Crop District 4 – Consul, Maple Creek and Leader areas)

Although recent warm weather has helped crops to advance in the southwest, many of them are suffering from a lack of moisture and will soon need rain to mature and fill. Rainfall this week ranged from trace amounts to 37 mm in the Cabri area. Since April 1, the Cadillac area has received 385 mm of rain.

Topsoil moisture conditions are deteriorating in many areas of the region, particularly the Maple Creek area as well as areas near the Alberta and United States borders. Topsoil moisture conditions on cropland are rated as 70 per cent adequate, 23 per cent short and seven per cent very short. Hay land and pasture topsoil moisture is rated as 60 per cent adequate, 28 per cent short and 12 per cent short. CD 4A is reporting that 40 per cent of both cropland and hay land and pasture are very short topsoil moisture, while CD 4B is reporting that 20 per cent of cropland and 23 per cent of hay land and pasture is very short of topsoil moisture.

Haying continues and livestock producers now have 61 per cent baled or put into silage, with an additional 18 per cent cut and ready for baling. Hay quality is rated as 10 per cent excellent, 77 per cent good, 10 per cent fair and three per cent poor.

Estimated Southwest Hay Yields (tons/acre) July 21, 2014		
	Dry land	**Irrigated Land**
Alfalfa	1.2	1.8
Brome/Alfalfa	1.1	1.4
Other Tame Hay	0.8	N/A
Wild Hay	0.7	N/A
Greenfeed	1.2	2.6

Most crop damage this week is attributed to drought conditions. Other sources include localized flooding, hail, wind, insects and diseases. Some producers are spraying for grasshoppers and diseases in later-seeded crops.

Farmers are busy haying, controlling pests, fixing fences and hauling grain.

East-Central Saskatchewan (Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas; Crop District 6A – Lumsden, Craik, Watrous and Clavet areas)

Crops continue to recover from recent flooding, although in some areas many fields have been significantly affected. Rain will be needed in many areas of the region to help remaining crops mature and fill. Rainfall this past week ranged from trace amounts to 30 mm in the Saskatoon area. The Foam Lake area has received the most rainfall since April 1 (506 mm).

Topsoil moisture conditions on cropland are rated as 15 per cent surplus, 82 per cent adequate and three per cent short. Hay land and pasture topsoil moisture is rated as nine per cent surplus, 86 per cent adequate and five per cent short. CD 5B is reporting that 22 per cent of cropland and 14 per cent of hay land and pasture have surplus topsoil moisture.

Estimated East-central Hay Yields (tons/acre) July 21, 2014		
	Dry land	**Irrigated Land**
Alfalfa	1.5	2.1
Brome/Alfalfa	1.4	2.1
Other Tame Hay	1.3	1
Wild Hay	1.1	N/A
Greenfeed	1.8	1.5

Livestock producers now have 27 per cent of the hay crop baled or put into silage, with an additional 26 per cent cut and ready for baling. At this time, hay quality is rated as 19 per cent excellent, 75 per cent good and six per cent fair.

Hail, wind and localized flooding have caused the most damage this week. Insects such as wheat midge and grasshoppers are also causing some damage. Producers are spraying for sclerotinia, fusarium and leaf spot diseases.

Farmers are busy haying and controlling diseases and insects.

West-Central Saskatchewan (Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas; Crop District 7A – Rosetown, Kindersley, Eston, Major; CD 7B - Kerrobert, Macklin, Wilkie and Biggar areas)

Storms moved through the region late last week, bringing high winds, baseball-sized hail and heavy rain. Many fields suffered extensive hail damage, lodging and localized flooding. There are also reports of destroyed buildings, damaged vehicles and toppled grain bins. Some areas received large amounts of rain within a couple of hours: the Biggar area received 82 mm. Since April 1, the Rosetown area has received 313 mm of rain, the most in the region.

Estimated West-central Hay Yields (tons/acre) July 21, 2014		
	Dry land	**Irrigated Land**
Alfalfa	1.4	2
Brome/Alfalfa	1.3	2.5
Other Tame Hay	1.0	2
Wild Hay	0.9	2.3
Greenfeed	1.7	3.2

Topsoil moisture conditions on cropland are rated as 15 per cent surplus, 79 per cent adequate and six per cent short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 81 per cent adequate and nine per cent short. CD 6B is reporting that 12 per cent of cropland and 14 per cent of hay land and pasture is short of topsoil moisture. Rain would be welcomed in many areas of the region to help crops mature and fill.

Haying is progressing and livestock producers now have 41 per cent of the hay crop baled or put into silage, with an additional 30 per cent cut and ready for baling. Hay quality is rated as nine per cent excellent, 83 per cent good and eight per cent fair.

The majority of crop damage this week is attributed to hail, wind and localized flooding. There are also reports of damage from grasshoppers and from diseases such as leaf spot and root rot.

Farmers are busy haying, hauling grain and controlling pests.

Northeastern Saskatchewan (Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas; Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas)

Some areas of the region received heavy rain last week that has flooded some fields and damaged crops. The Nipawin area reports receiving 84 mm of rain in one day, although there are reports of larger amounts near the Garrick area. Since April 1, the Arborfield area has received 345 mm of rain, the greatest amount for the region.

Estimated Northeast Hay Yields (tons/acre) July 21, 2014		
	Dry land	Irrigated Land
Alfalfa	1.5	N/A
Brome/Alfalfa	1.6	N/A
Other Tame Hay	1.0	N/A
Wild Hay	1.0	N/A
Greenfeed	1.0	N/A

Topsoil moisture conditions have worsened since last week and are now rated as 40 per cent surplus and 60 per cent adequate on cropland. Hay land and pasture topsoil moisture is rated as 67 per cent surplus and 33 per cent adequate. Many areas are in need of some warm and dry weather to dry up fields, while other areas will soon need a rain to help crops mature and fill.

Haying continues for livestock producers in the region as 35 per cent of the hay crop has been baled or put into silage. An additional 35 per cent is cut and ready for baling. Hay quality at this time is rated as eight per cent excellent, 67 per cent good, 17 per cent fair and eight per cent poor.

Crops are advancing quickly in the region, although wind and localized flooding have caused some damage. Producers are spraying for grasshoppers, wheat midge and diseases such as sclerotinia and leaf spots in later-seeded crops.

Farmers are busy haying, fixing fences and controlling pests.

Northwestern Saskatchewan (Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas; Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas)

Recent rain has helped crops advance, but it is slowing down haying. The Neilburg area is reporting that 38 mm of rain was received this past week, while the Hafford area is reporting 377 mm of rain since April 1, the greatest amount in the region.

Cropland topsoil moisture conditions are rated as three per cent surplus, 93 per cent adequate and four per cent short, while hay land and pasture topsoil moisture conditions are rated as one per cent surplus, 95 per cent adequate and four per cent short. For some in the region, rainfall would be welcomed, while for others, warm and dry weather is needed to help crops advance.

Thirty-six per cent of the hay crop has now been baled or put into silage, with an additional 26 per cent cut and ready for baling. At this time, hay quality is rated as 23 per cent excellent, 71 per cent good and six per cent fair.

Producers have been monitoring for insect pests and spraying for leaf disease and sclerotinia in later-seeded crops when weather allows. The majority of crop damage this past week was due to localized flooding, lodging and drought conditions in some areas.

Estimated Northwest Hay Yields (tons/acre) July 21, 2014		
	Dry land	**Irrigated Land**
Alfalfa	1.4	N/A
Brome/Alfalfa	1.5	N/A
Other Tame Hay	1.1	N/A
Wild Hay	1.0	N/A
Greenfeed	2.0	N/A

Farmers are busy haying, hauling grain and controlling pests.

SK (provincial) Crop Conditions - July 21, 2014							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	5	5	15	13	12	16	20
% good	59	49	63	67	59	59	45
% fair	25	43	18	16	25	19	17
% poor	10	2	3	4	3	5	12
% very poor	1	1	1	0	1	1	6
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	12	5	8	22	15	9
% good	57	56	78	66	58	55	81
% fair	27	23	16	21	15	21	10
% poor	6	8	1	5	4	4	0
% very poor	3	1	0	0	1	5	0
Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	4	20	12	14	8	14	22
% good	51	61	63	56	60	58	26
% fair	28	18	18	17	25	20	14
% poor	16	1	6	12	5	7	22
% very poor	1	0	1	1	2	1	16

	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	10	1	9	7	2	0
% good	47	59	71	64	45	32	60
% fair	29	22	25	22	27	33	40
% poor	10	8	2	5	12	13	0
% very poor	7	1	1	0	9	20	0
Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	8	12	3	16	3
% good	88	37	70	72	72	65	80
% fair	10	62	21	15	25	18	15
% poor	2	1	1	1	0	1	2
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	0	6	1	0	23	15	8
% good	78	80	85	60	57	65	82
% fair	22	13	13	40	15	18	10
% poor	0	1	1	0	3	2	0
% very poor	0	0	0	0	2	0	0
East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	9	6	9	15	5	7	8
% good	47	42	59	60	56	57	58
% fair	32	37	26	21	31	28	22
% poor	9	11	5	4	6	7	12
% very poor	3	4	1	0	2	1	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	9	5	3	0	16	14	0
% good	53	45	54	75	47	53	75
% fair	31	37	32	13	28	21	25
% poor	6	12	11	12	8	10	0
% very poor	1	1	0	0	1	2	0
West-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	16	7	24	16	22	24	35

% good	74	73	68	68	65	63	50
% fair	10	20	8	16	13	12	15
% poor	0	0	0	0	0	1	0
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	24	22	20	5	21	27	60
% good	68	66	63	75	71	56	30
% fair	8	12	17	20	8	16	10
% poor	0	0	0	0	0	1	0
% very poor	0	0	0	0	0	0	0
Northeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	11	0	11	10	0
% good	100	100	58	55	55	55	59
% fair	0	0	27	35	31	23	23
% poor	0	0	3	8	3	11	17
% very poor	0	0	1	2	0	1	1
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	9	0	5	1	0	N/A
% good	57	51	25	50	52	70	N/A
% fair	33	24	65	30	20	18	N/A
% poor	3	14	5	10	21	7	N/A
% very poor	0	2	5	5	6	5	N/A
Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	28	1	33	N/A	31	30	N/A
% good	37	99	57	N/A	62	63	N/A
% fair	35	0	10	N/A	7	7	N/A
% poor	0	0	0	N/A	0	0	N/A
% very poor	0	0	0	N/A	0	0	N/A
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	30	28	N/A	N/A	31	0	N/A
% good	70	56	N/A	N/A	51	75	N/A
% fair	0	15	N/A	N/A	17	25	N/A
% poor	0	1	N/A	N/A	1	0	N/A
% very poor	0	0	N/A	N/A	0	0	N/A

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending July 21, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 23, 2014

© 2014 Government of Saskatchewan

Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period July 15 to 21, 2014

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	5	233
	3	Enniskillen	5	186
	33	Moose Creek	N/A	244
	34	Browning	N/A	240
	61	Antler	N/A	432
	63	Moose Mountain	3	345
	64	Brock	3	245
	65	Tecumseh	5	329
1B	91	Maryfield	4	457
	122	Martin	NIL	600
	123	Silverwood	NIL	332
	124	Kingsley	NIL	320
	125 A	Chester	NIL	318
	125 B	Chester	NIL	342
	151 A	Rocanville	NIL	424
	151 B	Rocanville	NIL	503
	154	Elcapo	NIL	375
	155 A	Wolseley	0.5	304.5
	155 B	Wolseley	N/A	378
	67	Weyburn	2	198
2A	68	Brokenshell	NIL	200
	97	Wellington	NIL	214.5
2B	127 A	Francis	2	316
	127 B	Francis	0.3	225.7
	129	Bratt's Lake	4	239.5
	131 A	Baildon	1	284
	131 B	Baildon	10	362.2
	156 A	Indian Head	NIL	372.5
	156 B	Indian Head	NIL	408.5
	157	South Qu'Appelle	NIL	312
	160 A	Pense	NIL	299
	160 B	Pense	N/A	75
	161	Moose Jaw	4	372
	162	Caron	37	363
	191	Marquis	NIL	318
3ASE	38 A	Laurier	7	259.9
	38 B	Laurier	9	244
	39 A	The Gap	10	190
	39 B	The Gap	6	216
3ASW	10	Happy Valley	2	156
	12	Poplar Valley	N/A	256
	40 A	Bengough	3	191
	40 B	Bengough	NIL	224
	42	Willow Bunch	2	232
	43	Old Post	3	260
	70	Key West	NIL	63.6
	73 A	Stonehenge	3.8	246.3
	73 B	Stonehenge	8	316
	74	Wood River	NIL	175.5
3AN	101	Terrell	5	248
	102	Lake Johnston	20.3	215.4
	103	Sutton	NIL	150
	132 A	Hillsborough	5	223
	132 B	Hillsborough	6	291
	134	Shamrock	N/A	222
	193 A	Eyebrow	1	301
	193 B	Eyebrow	N/A	295
	224	Maple Bush	N/A	294
3BS	17	Val Marie	8	385
	75 A	Pinto Creek	NIL	248
	75 B	Pinto Creek	NIL	220
	76	Auvergne	N/A	185
	77	Wise Creek	5	288
	78	Grassy Creek	18	270
	105	Glenbain	1	179
	106	Whiska Creek	N/A	303
	107	Lac Pelletier	N/A	170
	108	Bone Creek	N/A	206
3BN	137	Swift Current	N/A	41.6
	138 A	Webb	17	221
	138 B	Webb	17.5	230.3
	166	Excelsior	9	207
	167	Sask. Landing	3.4	214.4
	168 A	Riverside	6	192
	168 B	Riverside	37.1	201.2
	226	Victory	10	227
	228	Lacadena	15.5	185
	257	Monet	13	138.6

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	49	White Valley	5	153
	51	Reno	18.4	115.7
	79	Arlington	9	264
	109 A	Carmichael	N/A	181
	109 B	Carmichael	N/A	149
	110	Piapot	NIL	135
	111	Maple Creek	NIL	93
4B	139	Gull Lake	6	207
	141	Big Stick	NIL	140
	142	Enterprise	NIL	184
	169	Pittville	N/A	162
	231	Happyland	27	175
5A	183 A	Fertile Belt	NIL	344
	183 B	Fertile Belt	NIL	369
	186	Abernethy	2	360
	211	Churchbridge	NIL	305
	213	Saltcoats	1	383
	216	Tullymet	N/A	272.5
	241	Calder	4	269
	243	Wallace	NIL	319
	244	Orkney	N/A	415
	245 A	Garry	2	400
	245 B	Garry	NIL	435
	245 C	Garry	2	432
	246	Ituna Bon Accord	2	371
	247	Kellross	3	335
	248	Touchwood	2	295
5B	271	Cote	1	277
	273	Sliding Hills	N/A	477
	277	Emerald	5	506
	305	Invermay	NIL	400
	307	Elfros	10	389
	308 A	Big Quill	10	292
	308 B	Big Quill	3	268
	331	Livingston	N/A	306
	334	Preeceville	N/A	112
	336	Sasman	3	293
	337	Lakeview	6	294
	338	Lakeside	13	313
	366	Kelvington	10	327
	367	Ponass Lake	25	314
6A	190 A	Dufferin	3	480
	190 B	Dufferin	9	386.5
	190 C	Dufferin	NIL	335
	190 D	Dufferin	1	189.5
	219	Longlaketon	1	301
	220	Mckillop	2	435
	221	Sarnia	2.9	381.7
	222	Craik	NIL	320
	251	Big Arm	6	414
	252	Arm River	1	370
	279	Mount Hope	4	330
	282	McCraney	5	362
	309	Prairie Rose	N/A	350
	310	Usborne	12	166.5
	312	Morris	10	113
	313	Lost River	N/A	306
	339	Leroy	13	380.1
	340	Wolverine	26	323
	341	Viscount	15	287
	343 A	Blucher	20	282.9
	343 B	Blucher	30	170
6B	254	Lorebum	N/A	246
	284	Rudy	17	298
	285	Fertile Valley	27	224
	286	Milden	N/A	153
	314	Dundurn	27	266
	344	Corman Park	32	295
	346	Perdue	31	283
	376	Eagle Creek	27	292
	403	Rosthern	34	254

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7A	259	Snipe Lake	N/A	20
	287	St. Andrews	31	313
	288	Pleasant Valley	27	257
	290 A	Kindersley	17	167
	290 B	Kindersley	N/A	105.5
	290 C	Kindersley	22	159.2
	292	Milton	29	218
	317 A	Marriott	N/A	127.5
	317 B	Marriott	81	278
	318	Mountain View	56	297
	320 A	Oakdale	29	197.1
	320 B	Oakdale	45	199
	321	Prairiedale	23	220
7B	347	Biggar	40	292
	350 A	Mariposa	82	211.6
	350 B	Mariposa	77	295
	351	Progress	75	240
	352	Heart's Hill	49	204
	377	Glenside	N/A	289
	378 A	Rosemount	55	245
	378 B	Rosemount	34	285
	379	Reford	53	311
	381	Grass Lake	N/A	206
	382	Eye Hill	62.5	231.6
	409	Buffalo	30	221
	410	Round Valley	40	285.4
8A	395	Porcupine	13	243
	397	Barrier Valley	13.1	304.3
	428	Star City	16	340
	456	Arborfield	40	345
	457	Connaught	33	265
	486	Moose Range	84	285
	487	Nipawin	N/A	214
8B	369	St. Peter	21	300
	370 A	Humboldt	28	331
	370 B	Humboldt	N/A	306
	371	Bayne	32	343
	372	Grant	22	328.5
	400	Three Lakes	12	342
	402	Fish Creek	20	315
	429	Flett's Springs	12	268
	459	Kinistino	10	308
	460	Birch Hills	13.7	282.9
9AE	488	Torch River	50	266
	491	Buckland	N/A	317
9AW	406	Mayfield	34	263
	435	Redberry	30	377
	436	Douglas	30	313
	463	Duck Lake	35	314.9
	467 A	Round Hill	30	362
9B	467 B	Round Hill	27	205
	438	Battle River	30	187
	440	Hillsdale	38	294.5
	442	Manitou Lake	34.3	216.1
	498 A	Parkdale	26	243.5
	498 B	Parkdale	27	200
	499	Mervin	35	191.5
	501 A	Frenchman Butte	27	268
	501 B	Frenchman Butte	30	214
	501 C	Frenchman Butte	31	259
	502	Britannia	27.5	184.5
	561	Loon Lake	35	277
	588 A	Meadow Lake	25	272
	588 B	Meadow Lake	20	303
	588 C	Meadow Lake	29	255
	588 D	Meadow Lake	13.8	321.2
	622	Beaver River	N/A	191.2

Municipality No: A, B, C and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2014
To: July 21, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200
Kilometers
Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 23, 2014

© 2014 Government of Saskatchewan

Cropland Topsoil Moisture Conditions

July 22, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 23, 2014

© 2014 Government of Saskatchewan

Hay and Pasture Topsoil Moisture Conditions

July 22, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 23, 2014

© 2014 Government of Saskatchewan